

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W.T.A. Harrison, J. Simpson and M. Weil

## The dicyclohexylamine salt of RG108 (*N*-phthalyl-*L*-tryptophan), a potential epigenetic modulator

Julie Braun, Irving Boittiaux, Anaëlle Tilborg, Didier Lambert and Johan Wouters

*Acta Cryst.* (2010). E66, o3175–o3176

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals Online is available from [journals.iucr.org](http://journals.iucr.org)

# The dicyclohexylamine salt of RG108 (*N*-phthalyl-L-tryptophan), a potential epigenetic modulator

Julie Braun,<sup>a</sup> Irving Boittiaux,<sup>a</sup> Anaelle Tilborg,<sup>a</sup> Didier Lambert<sup>b</sup> and Johan Wouters<sup>a\*</sup>

<sup>a</sup>Department of Chemistry, University of Namur, 61, Rue de Bruxelles, B-5000 Namur, Belgium, and <sup>b</sup>Louvain Drug Research Institute (LDRI), UCL, 50, Avenue Mounier, B-1200 Woluwe-Saint-Lambert, Belgium  
Correspondence e-mail: johan.wouters@fundp.ac.be


Received 5 October 2010; accepted 9 November 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å; disorder in solvent or counterion;  $R$  factor = 0.037;  $wR$  factor = 0.094; data-to-parameter ratio = 13.6.

The dicyclohexylamine salt of RG108 (*N*-phthalyl-L-tryptophan) co-crystallizes with a water molecule and a disordered molecule of dimethylformamide (DMF), *viz.* dicyclohexylaminium (*S*)-2-(1,3-dioxoisindolin-2-yl)-3-(1*H*-indol-3-yl)propanoate dimethylformamide solvate monohydrate,  $\text{C}_{12}\text{H}_{24}\text{N}^+ \cdot \text{C}_{19}\text{H}_{13}\text{N}_2\text{O}_4^- \cdot \text{C}_3\text{H}_7\text{NO} \cdot \text{H}_2\text{O}$ . The conformation of the deprotonated compound is constrained by charge-assisted strong hydrogen bonds with the dicyclohexylaminium ion and a dense hydrogen-bond network involving co-crystallized solvent molecules. The dihedral angle between the fused ring systems in the anion is  $58.35$  (4)°.

## Related literature

For the synthesis and biological evaluation, see: Brueckner *et al.* (2005).



## Experimental

### Crystal data

$\text{C}_{12}\text{H}_{24}\text{N}^+ \cdot \text{C}_{19}\text{H}_{13}\text{N}_2\text{O}_4^- \cdot \text{C}_3\text{H}_7\text{NO} \cdot \text{H}_2\text{O}$   
 $M_r = 606.75$   
Orthorhombic,  $P2_12_12_1$   
 $a = 9.0884$  (1) Å  
 $b = 15.0206$  (3) Å  
 $c = 24.4749$  (5) Å

$V = 3341.15$  (10) Å<sup>3</sup>  
 $Z = 4$   
Cu  $K\alpha$  radiation  
 $\mu = 0.67$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.55 \times 0.04 \times 0.03$  mm

### Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby (Gemini ultra Cu) detector  
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford)

Diffraction, 2009)  
 $T_{\min} = 0.967$ ,  $T_{\max} = 0.981$   
13362 measured reflections  
5552 independent reflections  
4936 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$   
 $wR(F^2) = 0.094$   
 $S = 1.01$   
5552 reflections  
409 parameters  
8 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.30$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>  
Absolute structure: Flack (1983), 2160 Friedel pairs  
Flack parameter: 0.02 (18)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N3}-\text{HN3B} \cdots \text{O4}^{\text{i}}$	0.86 (3)	1.88 (3)	2.7309 (19)	169 (2)
$\text{N2}-\text{H2} \cdots \text{O5}^{\text{ii}}$	0.86	1.97	2.813 (3)	165
$\text{N3}-\text{HN3A} \cdots \text{O3}^{\text{iii}}$	0.99 (3)	1.79 (3)	2.7740 (19)	173 (2)
$\text{O5}-\text{H5B} \cdots \text{O99}$	0.83 (4)	1.84 (4)	2.645 (4)	164 (4)
$\text{O5}-\text{H5C} \cdots \text{O1}^{\text{iv}}$	0.89 (4)	1.99 (4)	2.857 (2)	164 (3)

Symmetry codes: (i)  $x + 1, y, z$ ; (ii)  $x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$ ; (iii)  $x + \frac{1}{2}, -y + \frac{1}{2}, -z$ ; (iv)  $-x + \frac{1}{2}, -y, z + \frac{1}{2}$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported in part by the Fonds National de la Recherche Scientifique (FNRS, Belgium). Crystals were obtained during work for the Masters thesis of Miss Isabelle Bouhy. The authors thank Bernadette Norberg for her valuable help during the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2050).

## References

- Brueckner, B., Boy, R. G., Siedlecki, P., Munsch, T., Kliem, H. C., Zielenkiewicz, P., Suhai, S., Wiessler, M. & Lyko, F. (2005). *Cancer Res.* **65**, 6305–6311.  
Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.  
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.

Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## **supplementary materials**

*Acta Cryst.* (2010). E66, o3175-o3176 [ doi:10.1107/S160053681004626X ]

## The dicyclohexylamine salt of RG108 (*N*-phthalyl-L-tryptophan), a potential epigenetic modulator

J. Braun, I. Boittiaux, A. Tilborg, D. Lambert and J. Wouters

### Comment

RG108 (*N*-phthalyl-L-tryptophan) is a DNA methyltransferase (DMNT) inhibitor that was discovered by virtual screening (Brueckner *et al.*, 2005). It reactivates tumor suppressor gene expression in tumor cells by DNA demethylation. RG108 also inhibits human tumor cell line proliferation.

Isomer *S* (C9) of RG108 is obtained starting from L-tryptophan and phthalic anhydride in DMF.

The unprotonated carboxylate group (both C—O bond lengths are similar with C19—O3 = 1.246 (2) Å and C19—O4 = 1.247 (2) Å) of RG108 is close to the protonated *sp*<sup>3</sup> nitrogen atom (N3) of the amine (intermolecular O···N distances: O3···N3<sup>*i*</sup> = 2.774 (2) Å and O4···N3<sup>*ii*</sup> = 2.731 (2) Å; *i* = -1/2 + *x*, 1/2 - *y*, -*z*, *ii* = -1 + *x*, *y*, *z*, see also Table 1).

A water molecule (O5) has co-crystallized and is involved in the stability of the packing as it forms a network of H-bonds connecting the N—H (N2) of the indole ring of RG108 with a carbonyl function (O1) of the phthalimide ring of a symmetry-related molecule and the oxygen atom (O99) of a molecule of DMF solvent (Table 1).

In addition to H-bonding to the water, the extra (disordered) co-crystallized solvent molecule of DMF is tightly packed in a cavity formed by the aromatic heterocycles of RG108 (the phthalimide and the indole rings).

As a consequence of the dense packing (salt bridge, H-bonds and van der Waals interactions), the two aromatic, planar, heterocycles of RG108 are perpendicular (acute angle between the planes defined by the phthalimide and the indole rings = 58.35 (4)°).

### Experimental

Synthesis of the compound was made by micro-ave heating of L-tryptophane and phthalic anhydride in DMF by adapting the procedure described by Brueckner *et al.* (2005).

Crystals were obtained by evaporation at room temperature of a solution in mixture of methylene chloride and methanol (9/1).

### Refinement

The two H atoms of the water molecule and the two H atoms on (protonated) nitrogen N5 were located from  $\Delta F$  Fourier difference maps and their position refined. All other H atoms were placed at idealized positions and allowed to ride on their parent atoms.

## supplementary materials

Atoms of a DMF molecule were refined isotropically. Disorder has been taken into account by refining two sets of coordinates (0.7 and 0.3 occupancies respectively) for each atom of the DMF molecule. Bond lengths and valence angles were restrained to be similar in both disordered parts.

### Figures




Fig. 1. ORTEP view (with atom numbering) of the title compound. Only selected H atoms have been retained for clarity (on the chiral carbon, on the protonated nitrogen, and H involved in H-bonds). Displacement ellipsoids for non-H atoms are drawn at the 30% probability level.

### dicyclohexylaminium (S)-2-(1,3-dioxoisindolin-2-yl)-3-(1*H*-indol-3-yl)propanoate dimethylformamide solvate monohydrate

#### Crystal data

$C_{12}H_{24}N^+ \cdot C_{19}H_{13}N_2O_4^- \cdot C_3H_7NO \cdot H_2O$

$M_r = 606.75$

Orthorhombic,  $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 9.0884$  (1) Å

$b = 15.0206$  (3) Å

$c = 24.4749$  (5) Å

$V = 3341.15$  (10) Å<sup>3</sup>

$Z = 4$

$F(000) = 1304.0$

$D_x = 1.206$  Mg m<sup>-3</sup>

Cu  $K\alpha$  radiation,  $\lambda = 1.54184$  Å

Cell parameters from 7270 reflections

$\theta = 3.5\text{--}67.3^\circ$

$\mu = 0.67$  mm<sup>-1</sup>

$T = 293$  K

Needle, yellow

$0.55 \times 0.04 \times 0.03$  mm

#### Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby (Gemini ultra Cu) detector  
5552 independent reflections

Radiation source: Enhance Ultra (Cu) X-ray Source  
4936 reflections with  $I > 2\sigma(I)$

mirror  
Detector resolution: 10.3712 pixels mm<sup>-1</sup>

$R_{\text{int}} = 0.026$

$\theta_{\text{max}} = 67.4^\circ$ ,  $\theta_{\text{min}} = 3.5^\circ$

$\omega$  scans

$h = -10 \rightarrow 9$

Absorption correction: multi-scan  
(*CrysAlis PRO*; Oxford Diffraction, 2009)

$k = -15 \rightarrow 17$

$T_{\text{min}} = 0.967$ ,  $T_{\text{max}} = 0.981$

$l = -28 \rightarrow 29$

13362 measured reflections

#### Refinement

Refinement on  $F^2$

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.094$	$[1.00000 + 0.00000\exp(0.00(\sin\theta/\lambda)^2)] / [\sigma^2(F_o^2) + 0.0000 + 0.0000*P + (0.0611P)^2 + 0.0400\sin\theta/\lambda]$ where $P = 0.33333F_o^2 + 0.66667F_c^2$
$S = 1.01$	$(\Delta/\sigma)_{\max} < 0.001$
5552 reflections	$\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$
409 parameters	$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$
8 restraints	Absolute structure: Flack (1983), 2160 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.02 (18)

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.28198 (19)	0.08359 (12)	0.15901 (7)	0.0255 (4)	
C2	0.15124 (19)	0.04587 (13)	0.18771 (7)	0.0281 (4)	
C3	0.0899 (2)	-0.03769 (15)	0.18579 (8)	0.0371 (4)	
H3	0.1306	-0.0828	0.1646	0.044*	
C4	-0.0364 (2)	-0.05175 (17)	0.21717 (10)	0.0471 (6)	
H4	-0.0803	-0.1077	0.2171	0.056*	
C5	-0.0976 (2)	0.01558 (18)	0.24826 (9)	0.0466 (6)	
H5	-0.1820	0.0041	0.2686	0.056*	
C6	-0.0356 (2)	0.09991 (17)	0.24977 (9)	0.0409 (5)	
H6	-0.0774	0.1455	0.2703	0.049*	
C7	0.09121 (18)	0.11355 (14)	0.21952 (7)	0.0293 (4)	
C8	0.18476 (18)	0.19374 (14)	0.21386 (7)	0.0276 (4)	
C9	0.40737 (17)	0.23259 (12)	0.15742 (7)	0.0245 (4)	
H9	0.3868	0.2900	0.1749	0.029*	
C10	0.56294 (17)	0.20499 (14)	0.17540 (7)	0.0283 (4)	
H10A	0.5840	0.1463	0.1609	0.034*	
H10B	0.6334	0.2461	0.1596	0.034*	
C11	0.58403 (18)	0.20322 (13)	0.23613 (8)	0.0300 (4)	
C12	0.6295 (2)	0.27212 (15)	0.26820 (8)	0.0382 (5)	
H12	0.6514	0.3289	0.2553	0.046*	
C13	0.5969 (2)	0.15904 (15)	0.32564 (8)	0.0365 (4)	

## supplementary materials

---

C14	0.5908 (2)	0.10309 (17)	0.37118 (9)	0.0444 (5)
H14	0.6146	0.1238	0.4059	0.053*
C15	0.5484 (2)	0.01611 (17)	0.36273 (9)	0.0464 (6)
H15	0.5429	-0.0226	0.3923	0.056*
C16	0.5136 (2)	-0.01489 (15)	0.31049 (9)	0.0407 (5)
H16	0.4854	-0.0739	0.3059	0.049*
C17	0.52024 (19)	0.04037 (14)	0.26543 (8)	0.0329 (4)
H17	0.4971	0.0188	0.2309	0.039*
C18	0.56222 (18)	0.12900 (14)	0.27259 (7)	0.0297 (4)
C19	0.39345 (18)	0.24740 (12)	0.09508 (7)	0.0251 (4)
C20	1.0315 (2)	0.06650 (15)	0.02751 (8)	0.0362 (4)
H20A	1.1333	0.0726	0.0389	0.043*
H20B	1.0264	0.0797	-0.0112	0.043*
C21	0.9818 (3)	-0.02937 (15)	0.03696 (10)	0.0466 (5)
H21A	1.0377	-0.0688	0.0134	0.056*
H21B	1.0017	-0.0460	0.0745	0.056*
C22	0.8190 (3)	-0.04061 (16)	0.02532 (10)	0.0460 (5)
H22A	0.7894	-0.1011	0.0338	0.055*
H22B	0.8004	-0.0301	-0.0132	0.055*
C23	0.7290 (2)	0.02426 (15)	0.05937 (9)	0.0417 (5)
H23A	0.6254	0.0169	0.0510	0.050*
H23B	0.7431	0.0114	0.0979	0.050*
C24	0.7748 (2)	0.11973 (14)	0.04766 (8)	0.0325 (4)
H24A	0.7539	0.1338	0.0098	0.039*
H24B	0.7179	0.1598	0.0704	0.039*
C25	0.93764 (19)	0.13371 (13)	0.05872 (7)	0.0277 (4)
H25	0.9559	0.1267	0.0979	0.033*
C26	0.90346 (18)	0.30325 (13)	0.06517 (7)	0.0265 (4)
H26	0.7980	0.2957	0.0582	0.032*
C27	0.9557 (2)	0.38741 (14)	0.03688 (8)	0.0350 (4)
H27A	0.9352	0.3831	-0.0019	0.042*
H27B	1.0613	0.3931	0.0414	0.042*
C28	0.8803 (3)	0.46978 (15)	0.06000 (9)	0.0430 (5)
H28A	0.9198	0.5225	0.0425	0.052*
H28B	0.7758	0.4671	0.0521	0.052*
C29	0.9029 (2)	0.47659 (15)	0.12145 (9)	0.0424 (5)
H29A	1.0063	0.4862	0.1293	0.051*
H29B	0.8480	0.5270	0.1356	0.051*
C30	0.8515 (2)	0.39192 (16)	0.14926 (9)	0.0409 (5)
H30A	0.7461	0.3857	0.1443	0.049*
H30B	0.8706	0.3963	0.1882	0.049*
C31	0.92799 (19)	0.30959 (14)	0.12677 (7)	0.0317 (4)
H31A	0.8892	0.2568	0.1445	0.038*
H31B	1.0326	0.3129	0.1344	0.038*
N1	0.29621 (15)	0.17046 (10)	0.17706 (6)	0.0251 (3)
N2	0.6388 (2)	0.24668 (12)	0.32209 (7)	0.0429 (4)
H2	0.6661	0.2799	0.3489	0.051*
N3	0.98457 (16)	0.22521 (11)	0.04155 (6)	0.0269 (3)
O1	0.36330 (14)	0.04770 (9)	0.12646 (6)	0.0342 (3)



O2	0.17295 (15)	0.26576 (10)	0.23543 (6)	0.0374 (3)	
O3	0.50423 (13)	0.27765 (10)	0.07154 (5)	0.0331 (3)	
O4	0.27256 (13)	0.23090 (11)	0.07320 (6)	0.0383 (3)	
O5	0.1772 (2)	0.13211 (12)	0.59148 (8)	0.0539 (4)	
O99	0.0737 (4)	0.1983 (3)	0.49899 (14)	0.0750 (9)*	0.70
N99	0.1552 (5)	0.2310 (3)	0.41554 (16)	0.0510 (11)*	0.70
C88	0.1620 (4)	0.2142 (3)	0.46712 (15)	0.0517 (8)*	0.70
H88	0.2571	0.2153	0.4812	0.062*	0.70
C96	0.2644 (6)	0.2521 (4)	0.3781 (2)	0.0796 (13)*	0.70
H96A	0.2211	0.2610	0.3428	0.119*	0.70
H96B	0.3342	0.2042	0.3762	0.119*	0.70
H96C	0.3136	0.3056	0.3894	0.119*	0.70
C97	-0.0037 (4)	0.2278 (3)	0.39149 (17)	0.0656 (10)*	0.70
H97A	-0.0004	0.2411	0.3531	0.098*	0.70
H97B	-0.0642	0.2709	0.4098	0.098*	0.70
H97C	-0.0443	0.1694	0.3968	0.098*	0.70
O99B	0.0109 (6)	0.2045 (4)	0.5007 (2)	0.0357 (11)*	0.30
N99B	0.1149 (8)	0.2235 (5)	0.4188 (3)	0.0317 (18)*	0.30
C88B	0.0171 (12)	0.2275 (8)	0.4510 (4)	0.071 (3)*	0.30
H88B	-0.0696	0.2516	0.4373	0.085*	0.30
C96B	0.2691 (11)	0.2064 (7)	0.4520 (4)	0.067 (2)*	0.30
H96D	0.3491	0.2029	0.4265	0.100*	0.30
H96E	0.2624	0.1517	0.4721	0.100*	0.30
H96F	0.2860	0.2548	0.4769	0.100*	0.30
C97B	0.1649 (19)	0.2484 (12)	0.3692 (7)	0.112 (5)*	0.30
H97D	0.2612	0.2238	0.3633	0.168*	0.30
H97E	0.1701	0.3122	0.3674	0.168*	0.30
H97F	0.0992	0.2270	0.3414	0.168*	0.30
H5B	0.133 (4)	0.145 (3)	0.5631 (15)	0.072 (11)*	
H5C	0.150 (4)	0.076 (3)	0.5979 (13)	0.072 (10)*	
HN3B	1.077 (3)	0.2319 (16)	0.0481 (9)	0.033 (6)*	
HN3A	0.984 (3)	0.2272 (17)	0.0012 (11)	0.046 (6)*	

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0274 (8)	0.0247 (9)	0.0245 (8)	-0.0008 (8)	-0.0019 (7)	-0.0009 (7)
C2	0.0286 (8)	0.0303 (10)	0.0254 (8)	-0.0058 (8)	-0.0033 (7)	0.0066 (7)
C3	0.0407 (10)	0.0347 (11)	0.0358 (10)	-0.0087 (9)	-0.0072 (8)	0.0085 (8)
C4	0.0443 (11)	0.0470 (14)	0.0499 (12)	-0.0198 (11)	-0.0082 (10)	0.0180 (11)
C5	0.0300 (9)	0.0640 (16)	0.0456 (12)	-0.0128 (11)	0.0021 (8)	0.0228 (12)
C6	0.0299 (9)	0.0547 (14)	0.0382 (11)	-0.0004 (10)	0.0050 (8)	0.0111 (10)
C7	0.0259 (8)	0.0376 (11)	0.0242 (9)	-0.0010 (8)	-0.0007 (7)	0.0052 (8)
C8	0.0257 (8)	0.0340 (10)	0.0231 (8)	0.0020 (8)	0.0004 (6)	0.0010 (7)
C9	0.0256 (8)	0.0243 (9)	0.0235 (8)	-0.0044 (7)	0.0028 (6)	0.0002 (7)
C10	0.0239 (7)	0.0330 (10)	0.0279 (8)	-0.0048 (8)	-0.0010 (7)	0.0047 (8)
C11	0.0274 (8)	0.0323 (10)	0.0303 (9)	-0.0040 (8)	-0.0033 (7)	0.0067 (8)
C12	0.0496 (11)	0.0329 (11)	0.0320 (10)	-0.0066 (9)	-0.0117 (9)	0.0063 (8)

## supplementary materials

---

C13	0.0405 (9)	0.0386 (12)	0.0305 (9)	-0.0040 (9)	-0.0032 (8)	0.0047 (8)
C14	0.0525 (11)	0.0517 (14)	0.0291 (10)	-0.0029 (11)	-0.0046 (9)	0.0090 (10)
C15	0.0509 (12)	0.0486 (14)	0.0397 (11)	-0.0048 (11)	-0.0007 (9)	0.0212 (10)
C16	0.0394 (10)	0.0349 (12)	0.0479 (12)	-0.0061 (9)	-0.0029 (9)	0.0128 (9)
C17	0.0296 (8)	0.0345 (11)	0.0345 (10)	-0.0036 (8)	-0.0009 (7)	0.0025 (8)
C18	0.0266 (8)	0.0331 (11)	0.0293 (9)	-0.0016 (8)	-0.0010 (7)	0.0037 (8)
C19	0.0260 (8)	0.0242 (9)	0.0251 (8)	0.0023 (7)	-0.0001 (6)	0.0000 (7)
C20	0.0339 (9)	0.0391 (12)	0.0355 (10)	0.0120 (9)	-0.0027 (8)	-0.0033 (9)
C21	0.0586 (13)	0.0366 (13)	0.0445 (12)	0.0161 (11)	-0.0031 (10)	-0.0029 (10)
C22	0.0613 (13)	0.0301 (11)	0.0465 (12)	-0.0004 (11)	0.0006 (10)	0.0005 (9)
C23	0.0467 (11)	0.0334 (12)	0.0450 (12)	-0.0015 (10)	0.0041 (9)	0.0059 (9)
C24	0.0297 (8)	0.0311 (11)	0.0368 (10)	0.0033 (8)	0.0038 (7)	0.0022 (8)
C25	0.0309 (9)	0.0303 (10)	0.0219 (8)	0.0057 (8)	-0.0011 (7)	0.0008 (7)
C26	0.0237 (7)	0.0289 (9)	0.0268 (8)	0.0011 (7)	-0.0005 (6)	-0.0033 (7)
C27	0.0409 (10)	0.0352 (11)	0.0290 (9)	-0.0018 (9)	-0.0006 (8)	0.0002 (8)
C28	0.0516 (12)	0.0326 (11)	0.0448 (11)	-0.0035 (10)	-0.0019 (9)	-0.0012 (9)
C29	0.0456 (10)	0.0364 (12)	0.0452 (11)	-0.0059 (10)	0.0020 (9)	-0.0125 (10)
C30	0.0449 (10)	0.0424 (12)	0.0355 (11)	-0.0055 (10)	0.0090 (9)	-0.0106 (9)
C31	0.0295 (8)	0.0382 (11)	0.0274 (9)	-0.0010 (8)	0.0012 (7)	-0.0028 (8)
N1	0.0267 (7)	0.0236 (8)	0.0251 (7)	-0.0033 (6)	0.0035 (6)	-0.0021 (6)
N2	0.0647 (11)	0.0360 (10)	0.0280 (8)	-0.0077 (9)	-0.0123 (8)	0.0022 (7)
N3	0.0224 (7)	0.0340 (9)	0.0244 (8)	0.0040 (6)	-0.0023 (6)	-0.0013 (6)
O1	0.0384 (6)	0.0295 (7)	0.0346 (7)	-0.0011 (6)	0.0070 (6)	-0.0068 (6)
O2	0.0414 (7)	0.0334 (8)	0.0374 (7)	0.0041 (6)	0.0071 (6)	-0.0081 (6)
O3	0.0316 (6)	0.0428 (8)	0.0247 (6)	-0.0066 (6)	0.0013 (5)	0.0054 (6)
O4	0.0276 (6)	0.0523 (9)	0.0349 (7)	-0.0029 (6)	-0.0065 (5)	0.0039 (6)
O5	0.0860 (13)	0.0317 (9)	0.0442 (10)	0.0013 (9)	0.0080 (9)	0.0005 (7)

### *Geometric parameters (Å, °)*

C1—O1	1.213 (2)	C23—H23B	0.9700
C1—N1	1.384 (2)	C24—C25	1.519 (3)
C1—C2	1.492 (3)	C24—H24A	0.9700
C2—C3	1.374 (3)	C24—H24B	0.9700
C2—C7	1.392 (3)	C25—N3	1.499 (3)
C3—C4	1.397 (3)	C25—H25	0.9800
C3—H3	0.9300	C26—N3	1.500 (2)
C4—C5	1.383 (4)	C26—C27	1.517 (3)
C4—H4	0.9300	C26—C31	1.527 (2)
C5—C6	1.387 (4)	C26—H26	0.9800
C5—H5	0.9300	C27—C28	1.524 (3)
C6—C7	1.385 (3)	C27—H27A	0.9700
C6—H6	0.9300	C27—H27B	0.9700
C7—C8	1.481 (3)	C28—C29	1.522 (3)
C8—O2	1.209 (2)	C28—H28A	0.9700
C8—N1	1.400 (2)	C28—H28B	0.9700
C9—N1	1.457 (2)	C29—C30	1.516 (3)
C9—C10	1.538 (2)	C29—H29A	0.9700
C9—C19	1.547 (2)	C29—H29B	0.9700

C9—H9	0.9800	C30—C31	1.522 (3)
C10—C11	1.499 (3)	C30—H30A	0.9700
C10—H10A	0.9700	C30—H30B	0.9700
C10—H10B	0.9700	C31—H31A	0.9700
C11—C12	1.363 (3)	C31—H31B	0.9700
C11—C18	1.442 (3)	N2—H2	0.8600
C12—N2	1.376 (3)	N3—HN3B	0.86 (2)
C12—H12	0.9300	N3—HN3A	0.99 (3)
C13—N2	1.373 (3)	O5—H5B	0.82 (4)
C13—C14	1.397 (3)	O5—H5C	0.89 (4)
C13—C18	1.410 (3)	O99—C88	1.145 (5)
C14—C15	1.378 (4)	N99—C88	1.289 (5)
C14—H14	0.9300	N99—C96	1.387 (7)
C15—C16	1.397 (3)	N99—C97	1.561 (6)
C15—H15	0.9300	C88—H88	0.9300
C16—C17	1.381 (3)	C96—H96A	0.9600
C16—H16	0.9300	C96—H96B	0.9600
C17—C18	1.396 (3)	C96—H96C	0.9600
C17—H17	0.9300	C97—H97A	0.9600
C19—O3	1.246 (2)	C97—H97B	0.9600
C19—O4	1.247 (2)	C97—H97C	0.9600
C20—C25	1.527 (3)	O99B—C88B	1.266 (12)
C20—C21	1.527 (3)	N99B—C88B	1.189 (13)
C20—H20A	0.9700	N99B—C97B	1.350 (18)
C20—H20B	0.9700	N99B—C96B	1.640 (12)
C21—C22	1.517 (3)	C88B—H88B	0.9300
C21—H21A	0.9700	C96B—H96D	0.9600
C21—H21B	0.9700	C96B—H96E	0.9600
C22—C23	1.521 (3)	C96B—H96F	0.9600
C22—H22A	0.9700	C97B—H97D	0.9600
C22—H22B	0.9700	C97B—H97E	0.9600
C23—C24	1.521 (3)	C97B—H97F	0.9600
C23—H23A	0.9700		
O1—C1—N1	124.88 (17)	H24A—C24—H24B	108.0
O1—C1—C2	128.72 (18)	N3—C25—C24	110.73 (15)
N1—C1—C2	106.39 (15)	N3—C25—C20	107.89 (15)
C3—C2—C7	121.81 (17)	C24—C25—C20	111.34 (16)
C3—C2—C1	130.83 (19)	N3—C25—H25	108.9
C7—C2—C1	107.35 (16)	C24—C25—H25	108.9
C2—C3—C4	116.9 (2)	C20—C25—H25	108.9
C2—C3—H3	121.5	N3—C26—C27	108.74 (14)
C4—C3—H3	121.5	N3—C26—C31	110.93 (15)
C5—C4—C3	121.5 (2)	C27—C26—C31	110.66 (16)
C5—C4—H4	119.2	N3—C26—H26	108.8
C3—C4—H4	119.2	C27—C26—H26	108.8
C4—C5—C6	121.27 (19)	C31—C26—H26	108.8
C4—C5—H5	119.4	C26—C27—C28	111.49 (16)
C6—C5—H5	119.4	C26—C27—H27A	109.3
C7—C6—C5	117.3 (2)	C28—C27—H27A	109.3

## supplementary materials

---

C7—C6—H6	121.3	C26—C27—H27B	109.3
C5—C6—H6	121.3	C28—C27—H27B	109.3
C6—C7—C2	121.15 (19)	H27A—C27—H27B	108.0
C6—C7—C8	130.40 (19)	C29—C28—C27	111.16 (18)
C2—C7—C8	108.46 (15)	C29—C28—H28A	109.4
O2—C8—N1	124.68 (17)	C27—C28—H28A	109.4
O2—C8—C7	129.50 (17)	C29—C28—H28B	109.4
N1—C8—C7	105.82 (16)	C27—C28—H28B	109.4
N1—C9—C10	111.75 (15)	H28A—C28—H28B	108.0
N1—C9—C19	111.15 (14)	C30—C29—C28	110.23 (18)
C10—C9—C19	113.34 (13)	C30—C29—H29A	109.6
N1—C9—H9	106.7	C28—C29—H29A	109.6
C10—C9—H9	106.7	C30—C29—H29B	109.6
C19—C9—H9	106.7	C28—C29—H29B	109.6
C11—C10—C9	113.96 (14)	H29A—C29—H29B	108.1
C11—C10—H10A	108.8	C29—C30—C31	112.23 (16)
C9—C10—H10A	108.8	C29—C30—H30A	109.2
C11—C10—H10B	108.8	C31—C30—H30A	109.2
C9—C10—H10B	108.8	C29—C30—H30B	109.2
H10A—C10—H10B	107.7	C31—C30—H30B	109.2
C12—C11—C18	105.80 (16)	H30A—C30—H30B	107.9
C12—C11—C10	126.61 (18)	C30—C31—C26	109.94 (16)
C18—C11—C10	127.58 (18)	C30—C31—H31A	109.7
C11—C12—N2	111.09 (18)	C26—C31—H31A	109.7
C11—C12—H12	124.5	C30—C31—H31B	109.7
N2—C12—H12	124.5	C26—C31—H31B	109.7
N2—C13—C14	129.6 (2)	H31A—C31—H31B	108.2
N2—C13—C18	108.08 (17)	C1—N1—C8	111.92 (15)
C14—C13—C18	122.2 (2)	C1—N1—C9	124.30 (14)
C15—C14—C13	117.5 (2)	C8—N1—C9	123.65 (15)
C15—C14—H14	121.3	C13—N2—C12	108.05 (17)
C13—C14—H14	121.3	C13—N2—H2	126.0
C14—C15—C16	121.1 (2)	C12—N2—H2	126.0
C14—C15—H15	119.4	C25—N3—C26	117.94 (14)
C16—C15—H15	119.4	C25—N3—HN3B	109.4 (16)
C17—C16—C15	121.4 (2)	C26—N3—HN3B	108.4 (16)
C17—C16—H16	119.3	C25—N3—HN3A	107.8 (15)
C15—C16—H16	119.3	C26—N3—HN3A	110.9 (15)
C16—C17—C18	118.99 (19)	HN3B—N3—HN3A	101 (2)
C16—C17—H17	120.5	H5B—O5—H5C	103 (3)
C18—C17—H17	120.5	C88—N99—C96	131.1 (4)
C17—C18—C13	118.79 (18)	C88—N99—C97	114.1 (4)
C17—C18—C11	134.22 (18)	C96—N99—C97	114.8 (4)
C13—C18—C11	106.97 (17)	O99—C88—N99	132.4 (4)
O3—C19—O4	125.87 (17)	O99—C88—H88	113.8
O3—C19—C9	116.25 (15)	N99—C88—H88	113.8
O4—C19—C9	117.84 (15)	N99—C96—H96A	109.5
C25—C20—C21	112.50 (17)	N99—C96—H96B	109.5
C25—C20—H20A	109.1	H96A—C96—H96B	109.5

C21—C20—H20A	109.1	N99—C96—H96C	109.5
C25—C20—H20B	109.1	H96A—C96—H96C	109.5
C21—C20—H20B	109.1	H96B—C96—H96C	109.5
H20A—C20—H20B	107.8	N99—C97—H97A	109.5
C22—C21—C20	111.41 (18)	N99—C97—H97B	109.5
C22—C21—H21A	109.3	H97A—C97—H97B	109.5
C20—C21—H21A	109.3	N99—C97—H97C	109.5
C22—C21—H21B	109.3	H97A—C97—H97C	109.5
C20—C21—H21B	109.3	H97B—C97—H97C	109.5
H21A—C21—H21B	108.0	C88B—N99B—C97B	146.5 (11)
C21—C22—C23	110.53 (19)	C88B—N99B—C96B	108.5 (8)
C21—C22—H22A	109.5	C97B—N99B—C96B	101.6 (10)
C23—C22—H22A	109.5	N99B—C88B—O99B	131.1 (10)
C21—C22—H22B	109.5	N99B—C88B—H88B	114.5
C23—C22—H22B	109.5	O99B—C88B—H88B	114.5
H22A—C22—H22B	108.1	N99B—C96B—H96D	109.5
C24—C23—C22	110.70 (17)	N99B—C96B—H96E	109.5
C24—C23—H23A	109.5	H96D—C96B—H96E	109.5
C22—C23—H23A	109.5	N99B—C96B—H96F	109.5
C24—C23—H23B	109.5	H96D—C96B—H96F	109.5
C22—C23—H23B	109.5	H96E—C96B—H96F	109.5
H23A—C23—H23B	108.1	N99B—C97B—H97D	109.5
C25—C24—C23	111.33 (16)	N99B—C97B—H97E	109.5
C25—C24—H24A	109.4	H97D—C97B—H97E	109.5
C23—C24—H24A	109.4	N99B—C97B—H97F	109.5
C25—C24—H24B	109.4	H97D—C97B—H97F	109.5
C23—C24—H24B	109.4	H97E—C97B—H97F	109.5

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—HN3B...O4 <sup>i</sup>	0.86 (3)	1.88 (3)	2.7309 (19)	169 (2)
N2—H2...O5 <sup>ii</sup>	0.86	1.97	2.813 (3)	165
N3—HN3A...O3 <sup>iii</sup>	0.99 (3)	1.79 (3)	2.7740 (19)	173 (2)
O5—H5B...O99	0.83 (4)	1.84 (4)	2.645 (4)	164 (4)
O5—H5C...O1 <sup>iv</sup>	0.89 (4)	1.99 (4)	2.857 (2)	164 (3)

Symmetry codes: (i)  $x+1, y, z$ ; (ii)  $x+1/2, -y+1/2, -z+1$ ; (iii)  $x+1/2, -y+1/2, -z$ ; (iv)  $-x+1/2, -y, z+1/2$ .

Fig. 1

