

Proceedings

Open Access

Precision-cut liver slices in culture as a tool to assess the physiological involvement of Kupffer cells in hepatic metabolism

Audrey M Neyrinck*, Cristina Gomez and Nathalie M Delzenne

Address: Unité de Pharmacocinétique, Métabolisme, Nutrition et Toxicologie, Département des Sciences Pharmaceutiques, Université Catholique de Louvain, PMNT-UCL 73 avenue Mounier, B-1200 Brussels, Belgium

Email: Audrey M Neyrinck* - Audrey.Neyrinck@pmnt.ucl.ac.be; Cristina Gomez - Audrey.Neyrinck@pmnt.ucl.ac.be; Nathalie M Delzenne - Delzenne@pmnt.ucl.ac.be

* Corresponding author

from 11th International Symposium on the Cells of the Hepatic Sinusoid and their Relation to Other Cells
Tucson, Arizona, USA, 25–29 August, 2002

Published: 14 January 2004

Comparative Hepatology 2004, **3**(Suppl 1):S45

This article is available from: <http://www.comparative-hepatology.com/content/3/S1/S45>

Introduction

Hepatic macrophages have the capacity to secrete a tremendous array of molecules, which can be divided into 3 categories – cytokines (TNF- α), lipid mediators (prostaglandins PGE₂) and reactive intermediates (NO \cdot) – in response to stimulus, such as lipopolysaccharides (LPS) [1,2]. Such mediators are capable to modulate both the metabolism and the integrity of hepatocytes *in vitro* [2]. The physiological role of Kupffer cell in hepatic metabolism regulation has been approached in the present study by using the original *in vitro* model of precision-cut liver slices (PCLS) in culture; this model allows preserving the liver lobule architecture, by maintaining namely cell diversity in physiological proportion and cell-cell interactions [3]. First, we established whether non-parenchymal cells are still viable in rat PCLS and are able to respond to LPS *in vitro*; TNF- α , PGE₂, NO \cdot (reflecting NO \cdot release) were measured in the incubation medium of PCLS from rats previously treated with GdCl₃ – a specific inhibitor of Kupffer cell phagocytosis [4] – or NaCl as a control- in order to evaluate the contribution of Kupffer cell in mediator release. Moreover, by using the same model, we have investigated the role of Kupffer cell in the regulation of lipid synthesis in PCLS, in order to approach the biochemical mechanism explaining our last results, which indicate that the inhibition of Kupffer cell by GdCl₃ leads to triglycerides accumulation in liver tissue [5].

Methods

Materials

Male Wistar rats weighing 240–280 g were used for the preparation of PCLS or for isolation of hepatocytes. Most

chemicals of purest grade available were purchased from Sigma (Filter Service, Belgium), Roche Diagnostics Belgium or Invitrogen™ (Belgium). [1-¹⁴C]-acetic acid (specific activity 60 mCi/mmol) was obtained from Amersham Pharmacia Biotech Europe (Buckinghamshire, United Kingdom).

Study of mediator secretion by PCLS in culture

PCLS were prepared from treated with GdCl₃ (10 mg/kg *i.v.*) (Gd+) or NaCl 0.9% (Gd-) 24 h before liver removal according to a procedure previously described [6] and were incubated in William's E medium, supplemented with penicillin (100 IU/ml), streptomycin (100 micrograms/ml), glutamine (2 mM), insulin (100 nM) and bovine serum albumin 0.1 % containing LPS at 0 – 0.1 – 10 micrograms/ml. Medium was frozen after 2 h and 20 h of incubation for further analysis. PGE₂ and TNF- α concentration were measured in frozen aliquots incubation medium with immunoassay kits (PGE₂ Immunoassay, DE0100 and Quantikine rat TNF- α immunoassay, RTA00 from R&D Systems) whereas NO \cdot (NO₂⁻ + NO₃⁻) concentration was measured by the Griess reaction [7]. ATP content of PCLS was greater than 8 nmol/mg protein in all experiments.

Study of lipid synthesis by PCLS

PCLS were prepared from treated with GdCl₃ (10 mg/kg *i.v.*) (Gd+) or NaCl 0.9% (Gd-) 48 h before liver removal according to a procedure previously described [6]. PCLS were incubated as described above. Blood was collected from vena cava for serum PGE₂ measurement. After 2 h of preincubation, medium was frozen for further analysis

Table 1: Effect of Kupffer cell inhibition by GdCl₃ on lipid synthesis by PCLS

Acetate equivalent incorporated into lipids (nmol/mg prot.)	Gd-	Gd+
- phospholipids	3.7 ± 0.1	6.1 ± 0.6*
- triglycerides	2.4 ± 0.6	4.0 ± 0.8*
- cholesterol	0.8 ± 0.2	2.0 ± 0.2*

PCLS, prepared from 24 h-fasted rats pretreated with GdCl₃ (Gd+) or NaCl (Gd-) 2 days before the experiment, were incubated in medium containing 2 mM [¹⁴C]-acetate for 3 h. Values are means ± S.E.M (n ≥ 4) (*p < 0.05, Student t-test).

and PCLS were transferred into fresh medium containing 2 mM [¹⁴C]-acetate (0.2 mCi/mmol); after 3 h of incubation, PCLS were sonicated in 0.5 ml NaCl 0.05 M before lipid extraction and separation by thin-layer chromatography with hexane/ether/acetic acid (80:20:1) [8,9]. Spots corresponding to triglycerides, phospholipids and cholesterol were scrapped from the plate and counted in 10 ml scintillation fluid (Ultima Gold) in a beta counter.

Study of lipid synthesis by isolated hepatocytes in suspension

The hepatocytes were isolated from fed animals [10], through perfusion with a buffer containing Liberase™ (35 micrograms/ml). Cells were incubated in the presence PGE₂ (Cayman Chemicals) dissolved in DMSO at 10 micromolar and 2 mM [¹⁴C]-acetate (0.2 mCi/mmol) in the same medium described above. After 15 min of incubation, hepatocytes were pelleted and sonicated in 0.5 ml NaCl 0.05 M before lipid extraction according to Folch [8]. The chloroform phase was counted in 10 ml scintillation fluid (Ultima Gold) in a beta counter.

Statistical analysis

Values are presented as means ± S.E.M. In the study of mediator secretion by PCLS, statistical analysis was performed by two-way ANOVA with the Tukey's post hoc test (with SPSS™ statistical software). Other data were analysed by Student-*t* test. Statistical significance was set at p < 0.05.

Results

Validation of the PCLS model to assess the functionality of Kupffer cells

Unstimulated-PCLS in culture release in the medium significant amount of TNF-alpha, PGE₂ and NOx (Figure 1) which level increases with the time of incubation. TNF-alpha concentration increases significantly in the presence of LPS; the effect is dependent on the dose of LPS and is already present after 2 h of incubation. The extent of both PGE₂ and NOx production, is also dependent on LPS concentration; the increase in both parameters appears significantly only after 20 h of incubation. The administration of GdCl₃ 1 day before PCLS preparation strongly reduces


the basal production of the 3 mediators measured as well as upon stimulus by LPS.

Role of Kupffer cell in the physiological regulation of lipid synthesis in the liver tissue

PCLS obtained from rats treated with GdCl₃, 2 days before experiment, are characterised by a higher incorporation of [¹⁴C]-acetate into lipids (table 1), on one hand, and by a lower release of PGE₂ in the preincubation medium, on the other hand (PGE₂ concentration in the medium after 2 h: 5.4 ± 0.8 nM and 0.9 ± 0.1 nM for Gd- and Gd+ rats respectively; *p < 0.05, Student t-test). Moreover, PGE₂, measured in the serum obtained from vena cava, was much lower in animals treated with GdCl₃ 2 days before blood sampling (< 1 nM) as compared to control rats (26.7 ± 12.1 nM). The role of PGE₂ in the short-term control of lipid synthesis was assessed by incubation of isolated hepatocytes in suspension: the addition of PGE₂ (10 micromolar) to the culture medium rapidly decreases the incorporation of [¹⁴C]-acetate into total lipids in isolated hepatocytes after 15 minutes of incubation, total lipids – expressed in nmol acetate equivalent/mg protein – reached 1.3 ± 0.2 and only 0.87 ± 0.1 in control conditions (DMSO) and after addition of PGE₂ 10 micromolar, respectively (*p < 0.05, Student t-test).

Discussion

Our study demonstrates that GdCl₃, a specific inhibitor of Kupffer cell phagocytosis [4], decreases the rate of LPS-induced TNF-alpha, PGE₂ and NOx production by PCLS; those 3 molecules are known as typical mediators produced by macrophages (and namely by Kupffer cells in culture) upon stimulation by LPS (cytokines, lipid mediators and reactive intermediates) [10]. Moreover, we have shown that PCLS are able, independently of any stimulus (absence of LPS or any toxic agents) to produce TNF-alpha, NOx and PGE₂. In this case also, the release of mediators is decreased by previous GdCl₃ treatment. Therefore, we conclude that PCLS from GdCl₃-treated rats – compared to PCLS obtained from control rats – is a convenient in vitro system to study the complex Kupffer cell-hepatocyte interactions retained inside the liver tissue, in basal conditions or after an inflammatory stimulus. PGE₂

**Figure 1**

Effect of Kupffer cell inhibition on mediator secretion (TNF-alpha, PGE₂ and NOx) by PCLS, obtained from rats pretreated with GdCl₃ (Gd+) or NaCl (Gd-) 24 h before experiment; PCLS were incubated during 2 h and 20 h in the presence of LPS (from left to right, each group of three bars correspond to: 0 micrograms/ml, 0.1 micrograms/ml, 10 micrograms/ml). Values are means ± S.E.M (n ≥ 3; *p < 0.05 Gd+ vs Gd-, two-way ANOVA; ^ap < 0.05 LPS 10 micrograms/ml vs LPS 0 micrograms/ml and ^bp < 0.05 LPS 10 micrograms/ml vs LPS 0.1 micrograms/ml, two-way ANOVA followed by Tukey's post hoc test).

merits special attention: when PCLS are prepared from Gd+ animals, PGE₂ release is strongly depressed (mainly during short time incubation) suggesting that Kupffer cells are important producers of PGE₂ by the liver. Since GdCl₃ treatment leads to a strong depression of PGE₂ concentration in the serum, we may propose that Kupffer cells inside the liver constitute an important source of circulating PGE₂, known to exert pleiotropic effects outside the liver (control of lipolysis, platelet aggregation, gastric acid secretion, immunoregulation, neurotransmitter release, contraction-relaxation of smooth muscle) [11]. On the other hand, we show here that PGE₂ release by Kupffer cells, not only plays a role in the systemic availability of this prostaglandin, but is also able to exert paracrine effect on hepatocyte, with relevant metabolic consequences in the liver tissue: we had previously shown that, *in vivo*, the accumulation of lipids in the liver tissue after GdCl₃ administration, modifies the histological image of the liver, revealing steatosis [6]. We have demonstrated in the results presented here that the inhibition of Kupffer cells by GdCl₃ leads to a higher cholesterol, triglycerides and phospholipids synthesis by PCLS, a phenomenon related to a lower PGE₂ release. Since PGE₂ alone added in the culture medium of isolated hepatocytes decreases lipid synthesis, we may postulate that the lower PGE₂ secretion

after GdCl₃ treatment is, at least partly, responsible for a higher lipid synthesis inside the liver tissue. This is in favour of a role of Kupffer cell-released PGE₂ in the physiological control of lipid synthesis in hepatocytes. By which mechanism PGE₂ could control lipid synthesis in the liver tissue? The phosphorylation rate of hepatic enzymes involved in the lipogenesis or cholesterologenesis pathways (acetyl-CoA synthetase, HMG-CoA-reductase) could be modulated by PGE₂ which is released at concentration sufficient to alter the phosphorylation of specific proteins inside the hepatocytes [12]. The use of indomethacin, known to inhibit prostaglandin production, in the incubation medium of PCLS could help to elucidate the role of PGE₂ released by Kupffer cell in the regulation of hepatic metabolism.

References

1. Decker K: **Biologically active products of stimulated liver macrophages (Kupffer cells).** *Eur J Biochem* 1990, **192**:245-261.
2. Laskin DL, Pendino KJ: **Macrophages and inflammatory mediators in tissue injury.** *Annu Rev Pharmacol Toxicol* 1995, **35**:655-677.
3. C Lerche-Langrand, Toutain HJ: **Precision-cut liver slices: characteristics and use for in vitro pharmacotoxicology.** *Toxicology* 2000, **153**:221-253.
4. Hardonk MJ, Dijkhuis FWJ, Hulstaert CE, Koudstaal J: **Heterogeneity of rat liver and spleen macrophages in gadolinium chloride-induced elimination and repopulation.** *J Leukoc Biol* 1992, **52**:296-302.

5. Neyrinck AM, Taper HS, Delzenne NM: **Gadolinium chloride does not block the hypertriglyceridemia induced by LPS administration to rats.** In: *Cells of the Hepatic sinusoid Volume 8*. Edited by: Wisse E, Knook DL, de Zanger R, Fraser R. *Kupffer Cell Foundation, Leiden*; 2001:30-33.
6. Neyrinck AM, Taper HS, Gevers V, Declerck B, Delzenne NM: **Inhibition of Kupffer cell activity induces hepatic triglyceride synthesis in fasted rats, independent of lipopolysaccharide challenge.** *J Hepatol* 2002, **36**:466-473.
7. Bishop-Bailey D, Larkin SV, Warner TD, Chen G, Mitchell JA: **Characterization of the induction of nitric oxide synthase and cyclo-oxygenase in rat aorta in organ culture.** *Br J Pharmacol* 1997, **121**:125-133.
8. Folch J, Lee M, Sloane-Standley G: **A simple method for the isolation and purification of total lipids from animal tissues.** *J Biol Chem* 1957, **226**:497-509.
9. Liao W, Florèn CH: **Upregulation of low density lipoprotein receptor activity by TNF, a process of TNF-induced lipid synthesis and secretion.** *Lipids* 1994, **29**:679-684.
10. Roland CR, Naziruddin B, Mohanakumar T, Flye MW: **Gadolinium chloride inhibits Kupffer cell nitric oxide synthase (iNOS) induction.** *J Leukoc Biol* 1996, **60**:487-492.
11. Coleman RA, Smith VL, Narumiya S: **International union of pharmacology classification of prostanoid receptors: properties, distribution and structure of the receptors and their subtypes.** *Pharmacological Reviews* 1994, **46**:205-229.
12. Casteleijn E, Kuiper J, Van Rooij HC, Koster JF, Van Berckel TJ: **Conditioned media of Kupffer and endothelial cells influence protein phosphorylation in parenchymal liver cells. Involvement of prostaglandins.** *Biochem J* 1988, **252**:601-605.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

